

REVIEW ARTICLE

TRENDS in
Sport Sciences

2020; 27(3): 117-123

ISSN 2299-9590

DOI: 10.23829/TSS.2020.27.3-1

Physiological predictors of distance runners' performance: a narrative review

GERASIMOS GRIVAS

Abstract

The main purpose of this article was to review and discuss the current literature on the following physiological parameters: maximal oxygen uptake (VO_2max), running economy (RE), running velocity at vVO_2max (vVO_2max), time limit at vVO_2max (tlimit), running velocity at lactate threshold (vLT) and maximal speed (Vmax) on running performance. Many coaches and trainers believed that athletes with higher VO_2max have better performance, but a lot of studies have shown that VO_2max is a poor predictor of endurance performance. Alternatively, RE, vVO_2max , tlimit, vLT and Vmax were shown to be superior predictors of distance running performance. Therefore, if researchers and coaches can improve the values of all five parameters, the calculations suggested that the runner would be able to complete a marathon in 1:57:58. In conclusion, this review provides some practical suggestions as how to improve the performance in distance runners.

KEYWORDS: physiological parameters, performance, distance runners.

Received: 26 June 2020

Accepted: 22 August 2020

Corresponding author: grivasger@hotmail.com

University of Thessaly, Department of Physical Education and Sport Science, Trikala, Greece

Introduction

Many physiological variables are associated with aerobic function and are used to determine running performance [9]. These are: maximal oxygen uptake

(VO_2max), running economy (RE), running velocity at VO_2max (vVO_2max), time limit at vVO_2max (tlimit), running velocity at lactate threshold (vLT), and maximal speed (Vmax). The resulting question is which of these is the better predictor for performance in distance runners. It is very difficult to answer because studies use runners with different backgrounds and ability levels. It seems, however, that the most beneficial might be using the combination of several physiological factors to predict endurance performance in runners.

VO_2max is relatively homogeneous in elite runners, and within elite populations, race times were shown to have only a low to moderate correlation with VO_2max . For example, elite endurance athletes may have VO_2max values ranging between 70 and 85 $\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ [47]. Alternatively, physiological parameters like RE, vVO_2max , tlimit, vLT, and Vmax were shown to be superior predictors of distance running performance [1, 12, 20, 38]. The purpose of the current article is to discuss the role of VO_2max , RE, vVO_2max , tlimit, vLT, and Vmax and the significance of their implementation into distance runners' training. If the coaches or athletes consider the relevance of other physiological factors they will be able to maximize the running performance.

Maximal oxygen uptake (VO_2max)

VO_2max was first described by Hill and Lupton [28]. Over the following decades, VO_2max was accepted as a measure of the functional capacity of the cardiovascular system [27, 31, 46], and is taken as a measure of cardiorespiratory fitness. Average values in elite men long-distance runners range from 75 to 85 $\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$,

and from 60 to 75 ml·kg⁻¹·min⁻¹ in elite women long-distance runners [51, 54].

Paradoxically, there is a little relationship between VO₂max and race times among competitive long-distance runners [2]. A paradox is that a runner who improves VO₂max from 60 to 66 ml·kg⁻¹·min⁻¹ can usually be assured of an approximate 10% improvement in performance, but a runner with a VO₂max of 66 ml·kg⁻¹·min⁻¹ has no assurance that that is 10% better than a competitor with a VO₂max of 60 [2]. Studies have reported that VO₂max is a poor predictor of endurance performance when athletes of widely varying abilities are compared [34, 39, 41]. Legaz Arrese et al. [34] examined the relationship of VO₂max on performance in trained endurance runners. The results showed no relationship between changes in VO₂max and race performance. The study of Ramsbottom et al. [45] found that improvements in a 5 km trial were correlated with RE but not with VO₂max, while in the study by Paavolainen et al. [43] 5 km performance actually declined with improvements in VO₂max.

Training effects on VO₂max

Two of the most common endurance training strategies are interval training and continuous training methods [23]. The study of Pollock [44] has shown that improvement in VO₂max is directly related to intensity, duration, and frequency of training. The study by Helgerud et al. [27] revealed that the 15/15 (15-s intervals at 90-95% HRmax with 15 s of active resting periods) and the 4×4 min training (4×4-min interval training at 90-95% HRmax with 3 min of active resting periods at 70% HRmax between each interval) significantly increased the level of VO₂max in healthy students. However, long slow distance running (continuous run at 70% HRmax for 45 min) and lactate threshold running (continuous run at lactate threshold at 85% HRmax for 24.25 min) did not change their VO₂max. Another study [3] performed in young individuals with average cardiorespiratory fitness, shown that 20 sessions of periodized high volume interval training led to significant increases in VO₂max. Similarly, Gorostiaga et al. [24] when compared the effects of interval and continuous training programs on VO₂max showed that VO₂max was increased by 5% in the continuous training program, and 10% in the interval training program. The research clearly shows that continuous base training is an inefficient way to build an aerobic base and that the interval training improves VO₂max more effectively.

Running economy (RE)

Among the factors that may predict middle- and long-distance running performance, running economy (RE),

commonly defined as the steady-state VO₂ required at a given submaximal speed, has gathered the most attention over the last decade, although it is often still referred to as “being relatively ignored in the scientific literature” [4]. In the last decades, the researchers have focused on measuring RE in distance runners and many studies have found a strong association between RE and race performance. Some studies indicate that RE is an even better predictor of race performance among elite runners than VO₂max [16, 39, 47, 48].

Runners with good RE use less oxygen to run at a specific velocity compared to runners with less optimal economy [2]. When comparing two runners with similar VO₂max values the runner with better RE will achieve better performance time. Kenyan and Ethiopian runners have dominated middle- and long-distance running events compared to European runners. The runners from Africa do not have a higher VO₂max compared to European runners, but they have better performance [58]. The explanation is that African runners are typically smaller even compared to other elite runners, and studies have shown that smaller runners and runners with thinner and shorter lower limbs have better RE [14].

RE is a strong predictor for performance [32] because a runner with a greater economy will tend to work at lower percentages of VO₂max for various speeds than a runner who requires more oxygen and therefore has a poor economy [2]. For example, if two runners in a race have the same pace but one runner’s oxygen-consumption rate is 80% of VO₂max while the second runner’s oxygen-consumption rate is 90% of VO₂max, the first runner will continue the race for a greater distance than the second one. More specifically the first runner has greater RE than the second.

Numerous studies have examined the effects of RE in distance runners’ performance and found a strong association between RE and race performance. More specifically, Di Prampero et al. [18] reported that a 5% increase of RE induced a 3.8% increase in distance running performance. Weston et al. [57] investigated the RE and 10 km performance in African and Caucasian distance runners. African and Caucasian runners had similar race times in 10 km, but the African runners had a 13% lower VO₂max, but 5% better RE than Caucasians. This study indicates a greater RE and higher fractional utilization of VO₂peak in African distance runners. The study of Conley and Krahenbuhl [15] determined the relationship between RE and distance running performance in highly trained and experienced distance runners. All runners had similar VO₂max and within this elite cluster of finishers, 65.4% of the variation

observed in race performance time on the 10 km run could be explained by variation in RE.

On the other hand, only a few studies suggested that RE was not associated with running performance in competitive distance runners. Mooses et al. [39] suggested that in the homogenous group of Kenyan distance runners, RE can be compensated for by other factors (such as VO_2max) to maintain high-performance levels. Similar results are found in the study of Grant et al. [25] who reported that neither VO_2max nor RE was strongly correlated with the performance of 3 km. They also reported that vLT plays an important role in a 3 km running performance.

A variety of training strategies have been adopted in an attempt to improve RE. The most common training factors for improving RE are strength training (plyometric), tapering, hill, and pace-specific training (Figure 1).




Figure 1. Training factors for improving running economy

Strength training

RE improvements, as a consequence of strength training interventions, have been attributed to improved lower limb coordination and muscle coactivation, which would ultimately increase muscle stiffness and decrease ground contact times [35]. A meta-analysis published by Balsalobre-Fernández et al. [4] examined the effects of strength training on RE in highly trained distance runners. Authors reported that a strength training program including low to high-intensity resistance exercises and plyometric exercises performed 2-3 times per week for 8-12 weeks was an appropriate strategy to improve RE in the highly trained middle- and long-distance runners. The study by Li et al. [35], demonstrated improvement in RE and performance of 5 km running after complex training, which included a back squat + a drop jump from a 40 cm box; a Bulgarian split squat + a single leg hop; and a Romanian deadlift + a double leg 50 cm hurdle hop. Similarly, the study by Støren et al. [55] reported that maximal strength training for 8 weeks improved RE by 5.0% among well trained, long-distance runners, without a change in maximal oxygen uptake or body weight. Sedano et al [50] indicated that 12 weeks of combined and plyometric training lead

to improve RE. Paavolainen et al. [43] showed that 9 weeks of explosive-strength training improved RE (8%) and 5 km performance (3%) in moderately trained runners. The study by Saunders et al. [49] reported that in a group of highly-trained distance runners, 9 weeks of plyometric training improved RE by 4.1%, with likely mechanisms residing in the muscle, or alternatively by improving running mechanics. In a study of Blagrove et al. [13], ten weeks of strength training (2 days/week) added to the program of a postpubertal distance runner was highly likely to improve maximal speed and enhance RE by 3.2-3.7%. The study of Beattie et al. [7] demonstrated that 40 weeks of strength training can significantly improve maximal and reactive strength qualities, and RE by 5.0% in distance runners.

Tapering

Tapering is the reduction in training load before the competition or the final period before a major competition and it is very important for the athlete's performance. The aim of tapering is to maximize physiological adaptation with the reduction of accumulated fatigue [26]. Only 3 studies examined the effects of tapering on RE. The study of Houmard et al. [30] indicates that 7 days of tapered running improved distance running performance and RE by 6% in a group of well-trained endurance runners. In another study [52] RE at 60% vVO_2max was improved after 16 days of tapering. It is in agreement with the study of Houmard et al. [29] who investigated the effects of tapering on RE at 85% VO_2max speed after 2 and 3 weeks of taper and at 65% VO_2max after 3 weeks of taper.

Hill training

Hill training is another strategy that enhances RE. Only one study examined the effects of six-week hill training on RE in distance runners and showed its improvement by $2.4\% \pm 1.4\%$ [5].

Pace-specific training

Tempo runs are believed to enhance RE at the chosen training speed. Indeed, the study of Paavolainen et al. [43] suggested that muscle power may influence RE both at submaximal velocities and most probably at race pace. However, the study of Beneke and Hütler [8] indicated that training tends to improve economy the most at specific speeds used during training. This has important implications for the overall construction of a training plan. Moreover, 5 km runners should include a significant amount of training at the goal 5 km pace in order to optimize the economy at their desired intensity,

and marathoners should insert segments paced at their marathon goal speeds into their long runs and tempo runs [12].

Running velocity at VO_2 max (vVO_2 max)

vVO_2 max it is the minimum running velocity that elicits a runner's maximal rate of oxygen consumption or VO_2 max [2]. Some studies suggest that vVO_2 max is the best predictor of running performance [17]. vVO_2 max combines VO_2 max and RE into a single factor and explains differences in performance that VO_2 max or RE alone cannot [25]. The study of McLaughlin et al. [38] showed that among well-trained subjects – heterogeneous in VO_2 max and running performance – vVO_2 max is the best predictor of running performance because it integrates both maximal aerobic power and the RE. Morgan et al. [40] reported that there is a strong relationship between vVO_2 max and 10 km run time. Data from this study also suggest that vVO_2 max may be potentially useful as an index of training status and a sensitive, noninvasive predictor of distance running performance. McCormack et al. [37] reported that vVO_2 max was the best predictor of 3 km race performance in a group of collegiate distance runners with heterogeneous VO_2 max values. A study by Slattery et al. [53] showed similar results and explain that close relationship between vVO_2 max and 3 km race time may be due to the fact that middle distance races are completed at a velocity similar to that of VO_2 max, whereas velocities of longer distance races are closer to lactate threshold, and therefore vVO_2 max may become a better predictor of performance in the middle-distance events.

On the other hand, Grant et al. [25] found that vVO_2 max was the third-best predictor of 3 km run performance behind vLT and velocity at 4 mmol of blood lactate. According to Emerick et al. [19], this study failed to support the use of vVO_2 max as the best predictor of marathon performance in a group of recreational female runners. The study reinforced the notion that VO_2 max combined with weekly training distance elicits the best prediction of marathon performance. These findings suggest that recreational female runners should focus on increasing their VO_2 max and increasing their total training distance to improve their marathon performance. Endurance runners, coaches, and exercise physiologists gained the notion of why vVO_2 max is a much more useful performance predictor in distance running than VO_2 max. The latter contains no information about an athlete's RE. In case of a runner with high VO_2 max and low levels of RE, his performance could be disappointingly slow despite the high aerobic capacity [2].

Time limit at vVO_2 max

The time limit at vVO_2 max (tlimit) is also important for distance running performance and is a sister measurement of vVO_2 max. To determine the time limit at vVO_2 max the athlete runs at 100% of vVO_2 max until exhaustion, without slowing the pace or stopping. Direct measurement of tlimit indicates that it ranges from 150 s to 10 min and cannot last longer than 20-25 min [11, 33]. Billat [10] showed that the time limit at vVO_2 max was on average 6 min. Tlimit has a practical application in endurance athletes, for example, if two runners have similar values of vVO_2 max, the runner with higher tlimit will win the race.

Running velocity at lactate threshold (vLT)

Determining lactate threshold (LT), defined as the point at which blood lactate concentration increases exponentially with increasing exercise intensity, has been used to ascertain endurance capability, measure adaptations to training, and to predict performance potential [22]. Furthermore, LT is considered a valid performance indicator as there are strong linear correlations with endurance performance [21].

Running velocity at lactate threshold (vLT) is simply the velocity above which lactate begins to accumulate in the blood. Like RE and vVO_2 max, vLT is a strong physiological predictor of endurance performance [6]. Many coaches and researchers try to move the vLT to progressively faster speeds. Having a high vLT means that a runner can process pyruvate at greater rates and thus has the energy needed to run fast and long during endurance competitions [2].

In athletes with several years of training experience, VO_2 max may not improve any more, but vLT might increase by 3-10% depending on the chosen training program. It has been observed that individuals with similar VO_2 max have variability in endurance capacity and that highly trained athletes usually perform at a high percentage of their VO_2 max with minimum lactate accumulation [56, 59]. Furthermore, trained athletes accumulate less lactate than untrained athletes at a given submaximal workload.

Maximal speed (Vmax)

The study by Noakes et al. [42] was the first to report that maximal speed was a strong determinant of endurance performance. This study showed that peak treadmill velocity was the best laboratory predictor of running performance at 10, 21.1, 42.2, and 90 km distances in ultra-marathon runners, and it was also the best predictor of running performance for 10 and

21.1 km distances in marathon runners. The results of the Slattery et al. [53] study, showed that V_{max} was the single best predictor of 3-km running performance in experienced male triathletes and that both aerobic and anaerobic abilities are related to improved 3 km time trial performance. Since the assessment of V_{max} is relatively simple to implement, we suggest that determining V_{max} could be a practical method for monitoring performance changes in short-term endurance running events. The study of Paavolainen et al. [43] provides strong evidence that explosive training improves maximal running velocity and that the increases in maximal speed are closely coupled with improvements in endurance performance. This study examined 18 athletes who ran an all-out 20-meter sprint and a 5 km race (as fast as possible) on an indoor track. After 9 weeks the explosive-trained runners noted a 3% improvement and were 30 seconds faster in the 5 km event. Manoel et al. [36] compared the effects of 4 weeks of V_{max} or vVO_{2max} training in endurance runners. The results showed a significant effect of training on V_{max} and 10 km performance. It was concluded that V_{max} training promoted similar improvements as the training that included vVO_{2max} .

Conclusion

Many coaches and trainers believed that athletes with higher VO_{2max} have better performance. Many recent studies have shown that VO_{2max} alone is rather a poor predictor of endurance performance. It seems that actually, besides VO_{2max} , the physiological factors that are associated with aerobic capacity are RE, vVO_{2max} , t_{limit} , vLT , and V_{max} .

Conflicts of interest

The authors declare no conflict of interest.

Reference

- Allen WK, Seals DR, Hurley BF, Ehsani AA, Hagberg JM. Lactate threshold and distance-running performance in young and older endurance athletes. *J Appl Physiol.* 1985;58(4):1281-1284.
- Anderson O. *Running Science*. Champaign: Human Kinetics; 2013.
- Astorino TA, Edmunds RM, Clark A, King L, Gallant RM, Namm S, et al. Increased cardiac output and maximal oxygen uptake in response to ten sessions of high intensity interval training. *J Sports Med Phys Fitness.* 2018;58(1-2):164-171.
- Balsalobre-Fernández C, Santos-Concejero J, Grivas GV. The effects of strength training on running economy in highly trained runners: a systematic review with meta-analysis of controlled trials. *J Strength Cond Res.* 2016;30(8):2361-2368.
- Barnes KR, Hopkins WG, McGuigan MR, Kilding AE. Effects of different uphill interval-training programs on running economy and performance. *Int J Sports Physiol Perform.* 2013;8(6):639-647.
- Bassett DR Jr, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc.* 2000;32(1):70-84.
- Beattie K, Carson BP, Lyons M, Rossiter A, Kenny IC. The effect of strength training on performance indicators in distance runners. *J Strength Cond Res.* 2017;31(1):9-23.
- Beneke R, Hütler M. The effect of training on running economy and performance in recreational athletes. *Med Sci Sports Exerc.* 2005;37(10):1794-1799.
- Berg K. Endurance training and performance in runners: research limitations and unanswered questions. *Sports Med.* 2003;33(1):59-73.
- Billat VL, Flechet B, Petit B, Muriaux G, Koralsztejn JP. Interval training at VO_{2max} : effects on aerobic performance and overtraining markers. *Med Sci Sports Exerc.* 1999;31(1):156-163.
- Billat VL, Renoux JC, Pinoteau J, Petit B, Koralsztejn JP. Reproducibility of running time to exhaustion at VO_{2max} in subelite runners. *Med Sci Sports Exerc.* 1994;26(2):254-257.
- Bird SR, Theakston SC, Owen A, Nevill AM. Characteristics associated with 10-km running performance among a group of highly trained male endurance runners age 21-63 years. *J Aging Phys Act.* 2003;11(3):333-350.
- Blagrove RC, Howe LP, Cushion EJ, Spence A, Howatson G, Pedlar C, et al. Effects of strength training on postpubertal adolescent distance runners. *Med Sci Sports Exerc.* 2018;50(6):1224-1232.
- Borgen NT. Running performance, VO_{2max} , and running economy: the widespread issue of endogenous selection bias. *Sports Med.* 2018;48(5):1049-1058.
- Conley DL, Krahenbuhl GS. Running economy and distance running performance of highly trained athletes. *Med Sci Sports Exerc.* 1980;12(5):357-360.
- Costill DL, Thomason H, Roberts E. Fractional utilization of the aerobic capacity during distance running. *Med Sci Sports.* 1973;5(4):248-252.
- Demarie S, Koralsztejn JP, Billat V. Time limit and time at VO_{2max} during a continuous and an intermittent run. *J Sports Med Phys Fitness.* 2000;40(2):96-102.
- Di Prampero PE, Capelli C, Pagliaro P, Antonutto G, Girardis M, Zamparo P, et al. Energetics of best performances in middle-distance running. *J Appl Physiol.* 1993;74(5):2318-2324.

19. Emerick P, Teed K, Rusk G, Fernhall B. Predictors of marathon performance in female runners. *Sports Med Train Rehabil.* 1997;8(1):23-36.
20. Farrell PA, Wilmore JH, Coyle EF, Billing JE, Costill DL. Plasma lactate accumulation and distance running performance. *Med Sci Sports.* 1979;11(4):338-344.
21. Faude O, Kindermann W, Meyer T. Lactate threshold concepts: how valid are they? *Sports Med.* 2009; 39(6): 469-490.
22. Forsyth J, Burt D, Ridley F, Mann C. Using lactate threshold to predict 5-km treadmill running performance in veteran athletes. *Biol Sport.* 2017;34(3):233-237.
23. González-Mohíno F, Santos-Concejero J, Yustres I, González-Ravé JM. The effects of interval and continuous training on the oxygen cost of running in recreational runners: a systematic review and meta-analysis. *Sports Med.* 2020;50(2):283-294.
24. Gorostiaga EM, Walter CB, Foster C, Hickson RC. Uniqueness of interval and continuous training at the same maintained exercise intensity. *Eur J Appl Physiol Occup Physiol.* 1991;63(2):101-107.
25. Grant S, Craig I, Wilson J, Aitchison T. The relationship between 3 km running performance and selected physiological variables. *J Sports Sci.* 1997;15(4):403-410.
26. Grivas GV. The effects of tapering on performance in elite endurance runners: a systematic review. *Int J Sports Sci.* 2018;8(1):8-13.
27. Helgerud J, Høydal K, Wang E, Karlsen T, Berg P, Bjerkaas M, et al. Aerobic high-intensity intervals improve VO_2 max more than moderate training. *Med Sci Sports Exerc.* 2007;39(4):665-671.
28. Hill AV, Lupton H. Muscular exercise, lactic acid, and the supply and utilization of oxygen. *QJM.* 1924; 16(62):135-171.
29. Houmard JA, Costill DL, Mitchell JB, Park SH, Hickner RC, Roemmich JN. Reduced training maintains performance in distance runners. *Int J Sports Med.* 1990;11(1):46-52.
30. Houmard JA, Scott BK, Justice CL, Chenier TC. The effects of taper on performance in distance runners. *Med Sci Sports Exerc.* 1994;26(5):624-631.
31. Hutchinson A. *Endure: Mind, Body, and the Curiously Elastic Limits of Human Performance.* HarperCollins Publishers; 2018.
32. Joyner MJ. Modeling: optimal marathon performance on the basis of physiological factors. *J Appl Physiol.* 1991;70(2):683-687.
33. Lacour JR, Padilla-Magunacelaya S, Chatard JC, Arzac L, Barthélémy JC. Assessment of running velocity at maximal oxygen uptake. *Eur J Appl Physiol Occup Physiol.* 1991; 62(2):77-82.
34. Legaz Arrese A, Serrano Ostáriz E, Jcasajús Mallén JA, Munguía Izquierdo D. The changes in running performance and maximal oxygen uptake after long-term training in elite athletes. *J Sports Med Phys Fitness.* 2005;45(4): 435-440.
35. Li F, Wang R, Newton RU, Sutton D, Shi Y, Ding H. Effects of complex training versus heavy resistance training on neuromuscular adaptation, running economy and 5-km performance in well-trained distance runners. *Peer J.* 2019 Apr 7:e6787.
36. Manoel FA, da Silva DF, Lima JRP, Machado FA. Peak velocity and its time limit are as good as the velocity associated with VO_2 max for training prescription in runners. *Sports Med Int Open.* 2017 Jan 31(1):E8-E15.
37. McCormack W, Shoepe T, Almstedt H, Jennings C, Capel L. Velocity at maximal oxygen uptake best predicts 3 km race time in collegiate distance runners. *J Hum Sport Exerc.* 2018;13(3):631-638.
38. McLaughlin JE, Howley ET, Bassett DR Jr, Thompson DL, Fitzhugh EC. Test of the classic model for predicting endurance running performance. *Med Sci Sports Exerc.* 2010;42(5):991-997.
39. Mooses M, Mooses K, Haile DW, Durussel J, Kaasik P, Pitsiladis YP. Dissociation between running economy and running performance in elite Kenyan distance runners. *J Sports Sci.* 2015;33(2):136-144.
40. Morgan DW, Baldini FD, Martin PE, Kohrt WM. Ten kilometer performance and predicted velocity at VO_2 max among well-trained male runners. *Med Sci Sports Exerc.* 1989;21(1):78-83.
41. Morgan D, Martin P, Craib M, Caruso C, Clifton R, Hopewell R. Effect of step length optimization on the aerobic demand of running. *J Appl Physiol.* 1994;77(1):245-251.
42. Noakes TD, Myburgh KH, Schall R. Peak treadmill running velocity during the VO_2 max test predicts running performance. *J Sports Sci.* 1990;8(1):35-45.
43. Paavolainen L, Häkkinen K, Hämmäläinen I, Nummela A, Rusko H. Explosive-strength training improves 5-km running time by improving running economy and muscle power. *J Appl Physiol.* 1999;86(5):1527-1533.
44. Pollock ML. Submaximal and maximal working capacity of elite distance runners. Part I: Cardiorespiratory aspects. *Ann N Y Acad Sci.* 1977;301:310-322.
45. Ramsbottom R, Williams C, Fleming N, Nute ML. Training induced physiological and metabolic changes associated with improvements in running performance. *Br J Sports Med.* 1989;23(3):171-176.
46. Rowell LB. Human cardiovascular adjustments to exercise and thermal stress. *Physiol Rev.* 1974;54(1):75-159.
47. Saltin B, Astrand PO. Maximal oxygen uptake in athletes. *J Appl Physiol.* 1967;23(3):353-358.

48. Saunders PU, Pyne DB, Telford RD, Hawley JA. Factors affecting running economy in trained distance runners. *Sports Med.* 2004;34(7):465-485.
49. Saunders PU, Telford RD, Pyne DB, Peltola EM, Cunningham RB, Gore CJ, et al. Short-term plyometric training improves running economy in highly trained middle and long distance runners. *J Strength Cond Res.* 2006;20(4):947-954.
50. Sedano S, Marín PJ, Cuadrado G, Redondo JC. Concurrent training in elite male runners: the influence of strength versus muscular endurance training on performance outcomes. *J Strength Cond Res.* 2013;27(9):2433-2443.
51. Sjödín B, Svedenhag J. Applied physiology of marathon running. *Sports Med.* 1985;2(2):83-99.
52. Skovgaard C, Almquist NW, Kvorning T, Christensen PM, Bangsbo J. Effect of tapering after a period of high-volume sprint interval training on running performance and muscular adaptations in moderately trained runners. *J Appl Physiol.* 2018;124(2):259-267.
53. Slattery KM, Wallace LK, Murphy AJ, Coutts AJ. Physiological determinants of three-kilometer running performance in experienced triathletes. *J Strength Cond Res.* 2006;20(1):47-52.
54. Smith D, Telford R, Peltola E, Tumilty D. Protocols for the physiological assessment of high-performance runners. In: Gore CJ, editor. *Physiological Tests for Elite Athletes*, Australian Sports Commission. Human Kinetics: Champaign IL; 2000. pp. 334-344.
55. Støren O, Helgerud J, Støa EM, Hoff J. Maximal strength training improves running economy in distance runners. *Med Sci Sports Exerc.* 2008;40(6):1087-1092.
56. Tanaka K, Matsuura Y, Matsuzaka A, Hirakoba K, Kumagai S, Sun SO, et al. A longitudinal assessment of anaerobic threshold and distance-running performance. *Med Sci Sports Exerc.* 1984;16(3):278-282.
57. Weston AR, Mbambo Z, Myburgh KH. Running economy of African and Caucasian distance runners. *Med Sci Sports Exerc.* 2000;32(6):1130-1134.
58. Wilber RL, Pitsiladis YP. Kenyan and Ethiopian distance runners: what makes them so good? *Int J Sports Physiol Perform.* 2012;7(2):92-102.
59. Withers RT, Sherman WM, Miller JM, Costill DL. Specificity of the anaerobic threshold in endurance trained cyclists and runners. *Eur J Appl Physiol Occup Physiol.* 1981;47(1):93-104.